



Global distribution of *Strongyloides stercoralis*. Black = hyperendemic (>5%), high density dots = endemic (1-5%), sparse dots = sporadic (<1%). Overseas distribution: Stutchler (1981), in: Grove Grove DI. Strongyloidiasis. A major roundworm of man. 1989. Taylor & Francis Ltd., U.K. & U.S.A.
Australian distribution compiled from various studies (see below).

LIST OF PLACES IN AUSTRALIA WHERE STRONGYLOIDES DATA ARE AVAILABLE

Donydji 1986	29	41%	Stool	Flannery & White 1993 (1)
Yirrkala 1986	185	60%	IgG	Flannery & White 1993 (1)
Galiwin'ku	200	15%	Harada&Mori	Aland 1996 (2)
Croker Island	150	40%	IgG	Van Ingen 2003 (3)
Miwatj	508	35%	IgG	Page 2004 (4)
<i>Children under 15 years</i>				
Doomadgee 1974-90	1488	27% (12%)	Stool	Prociv & Luke 1993 (5)
Aurukun 1973-90	938	6.5%	Stool	Prociv & Luke 1993
Gununa 1978-91	928	5.6%	Stool	Prociv & Luke 1993
Weipa 1974-88	859	5%	Stool	Prociv & Luke 1993
Yarrabah 1975-86	1465	3%	Stool	Prociv & Luke 1993
Kowamyama 1974-87	1120	3%	Stool	Prociv & Luke 1993
Lockhart River 1973-91	1568	3%	Stool	Prociv & Luke 1993
Mt Isa 1975-89	511	2%	Stool	Prociv & Luke 1993
Cherbourg 1974-89	2462	2%	Stool	Prociv & Luke 1993
Woorabinda 2005	715	15%	IgG	Lord 2005
Billiluna 1990	98	37%	IgG	Sampson 2003 (6)
Kalumburu 1999	229	11.4%	IgG	Sampson 2003
<i>Children aged from 3 to 18 years</i>				
Kalumburu 1986	44	5%	IgG	Sampson 2003
One Arm Point 1986	67	4.5%	IgG	Sampson 2003
Loona 1986	57	58%	IgG	Sampson 2003
Lombardina 1986	30	10%	IgG	Sampson 2003
Beagle Bay 1986	28	17.8%	IgG	Sampson 2003

References

Aland K. 1996 Worm project at Galiwin'ku. Working Together 6 (6): 10.

See also conference proceedings available at: <http://www.sph.uq.edu.au/ACITHN/conf97/papers97/indigenoushealth.htm>

Barkworth J, Tye V, Barry M, Baggs L, Miller L, Smith S, Young M, Murphy K, Rewquandie D, Mick A, Doolan F. 2005 Woorabinda *Strongyloides* Eradication Project 2004-2007. Second National Workshop on Strongyloidiasis, Yeppoon 11 June 2005.

Fisher D, McCarry F, Currie B. 1993 Strongyloidiasis in the Northern Territory. "Under-recognised and under-treated?" Med. J. Aust. 159:88-90.

Flannery G, White N. 1993 Immunological parameters in northeast Arnhem Land Aborigines: consequences of changing settlement and lifestyles. In Urban Ecology and Health in the Third World. Ed. L.M. Schell, M.T. Smith & A. Bilsborough, pp. 202-220. Cambridge Uni. Press.

Lord R. 2005 *Strongyloides* serology from Woorabinda. Third National Strongyloidiasis Workshop Yeppoon 10-11 June 2005.

Page W. 2004 Report on Miwatj Strongyloidiasis Study. Implementing best practice in the eradication of chronic strongyloidiasis for clients of Miwatj Health Aboriginal Corporation. Report to ACCRM Darwin.

Prociv P, Russell L. 1993 Observations on strongyloidiasis in Queensland Aboriginal communities. Medical Journal of Australia 158: 160-163.

Sampson I, Smith DW, MacKenzie B. 2003 Serological diagnosis of *Strongyloides stercoralis* infection. Second National Strongyloidiasis Workshop, Brisbane 2003, URL. www.jcu.edu.au/school/phtm/PHTM/ss/

Van Ingen I. 2003 Strongyloidiasis in an island community. Progress of a treatment programme. Second National Strongyloidiasis Workshop, Brisbane 2003, URL. www.jcu.edu.au/school/phtm/PHTM/ss/